

ARQ-151, Roflumilast Cream, Improved Chronic Plaque Psoriasis in Phase 2b Study

Mark G. Lebwohl¹, Kim A. Papp², Linda Stein Gold³, Melinda J. Gooderham⁴, Leon H. Kircik⁵, Zoe D. Draelos⁶, Steven E. Kempers⁷, Mathew Zirwas⁸, Kathleen Smith⁹, David W. Osborne⁹, Marie-Louise Trotman¹⁰, Lynn Navale⁹, Charlotte Merritt⁹, David R. Berk⁹, Howard Welgus⁹

¹Icahn School of Medicine at Mount Sinai, New York, NY, USA; ²Probity Medical Research and K Papp Clinical Research, Waterloo, ON, Canada; ³Henry Ford Medical Center, Detroit, MI, USA; ⁴SkiN Centre for Dermatology, Probity Medical Research and Queen's University, Peterborough, ON, Canada; ⁵Icahn School of Medicine, Mount Sinai, New York, NY, and Skin Sciences, Louisville, KY, USA; ⁶Dermatology Consulting Services, High Point, NC, USA; ⁷Minnesota Clinical Study Center, Fridley, MN, USA; ⁸Probity Medical Research and Dermatologists of the Central States, Bexley, OH, USA; ⁹Arcutis, Inc., Westlake Village, CA, USA; ¹⁰ML Trotman Consulting, LLC, Newbury Park, CA, USA.

Disclosures

L. Stein Gold is an investigator for AbbVie, Arcutis, Celgene, Dermavant, Eli Lilly, LEO Pharma, Novartis, and Ortho Dermatologics; serves as an advisor for Celgene, Dermavant, LEO Pharma, Novartis, and Ortho Dermatologics; and is a speaker for LEO Pharma and Ortho Dermatologics.

M.G. Lebwohl reports receipt of research funds from AbbVie, Amgen, Arcutis, AstraZeneca, Boehringer Ingelheim, Celgene, Clinuvel, Eli Lilly, Incyte, Janssen Research & Development, LLC, Kadmon Corp., LLC, LEO Pharma, Medimmune, Novartis, Ortho Dermatologics, Pfizer, SCIderm, UCB, Inc., and Vidac, and serves as a consultant for Allergan, Almirall, Arcutis, Avotres Therapeutics, BirchBioMed Inc., Boehringer Ingelheim, Bristol-Myers Squibb, Cara Therapeutics, Castle Biosciences, Corrona, Dermavant Sciences, EMD Serono, Evelo Biosciences, Foundation for Research and Education in Dermatology, Inozyme Pharma, LEO Pharma, Meiji Seika Pharma, Menlo, Mitsubishi Pharma, Neuroderm, Pfizer, Promius/Dr. Reddy's Laboratories, Theravance, and Verrica. **K.A. Papp** is an investigator, consultant, speaker, and/or scientific officer and/or has served on steering committees or advisory boards for AbbVie, Akros, Amgen, Anacor, Arcutis, Astellas, Bausch Health/Valeant, Baxalta, Boehringer Ingelheim, Bristol-Myers Squibb, Can-Fite BioPharma, Celgene, Coherus, Dermira, Dow Pharma, Eli Lilly, Galderma, Genentech, Gilead, GSK, InflaRx GmbH, Janssen, Kyowa Hakko Kirin, LEO Pharma, Medimmune, Meiji Seika Pharma, Merck (MSD), Merck-Serono, Mitsubishi Pharma, Moberg Pharma, Novartis, Pfizer, PRCL Research, Regeneron, Roche, Sanofi-Aventis/Genzyme, Sun Pharma, Takeda, and UCB. **M.J. Gooderham** has been a speaker, advisory board member, investigator, and/or consultant for AbbVie, Amgen, Akros, Arcutis, Boehringer Ingelheim, BMS, Celgene, Dermira, Eli Lilly, Galderma, GSK, Janssen, Kyowa Kirin, LEO Pharma, Medimmune, Merck, Novartis, Pfizer, Regeneron, Sanofi Genzyme, UCB, and Valeant/Bausch. **L.H. Kircik** is an investigator, consultant, speaker, and/or serves on advisory boards for Abbott Laboratories, Acambis, Aclaris, Allergan, Inc., Almirall, Amgen Inc., Anacor Pharmaceuticals, Assos Pharma, Astellas Pharma US, Inc., Asubio, Berlex Laboratories (Bayer HealthCare Pharmaceuticals), Biogen-Idec, Biolife, Biopelle, Boehringer Ingelheim, Breckinridge Pharma, Celgene, Cellceutix, Centocor, Cipher, Coherus, Colbar, CollaGenex, Combinatrix, Connetics Corporation, Coria, Dermavant, Dermik Laboratories, Dermira, Dow Pharmaceutical Sciences, Inc., Dusa, Eli Lilly, Embil Pharmaceuticals, EOS, Exeltis, Ferndale Laboratories, Inc., Foamix, Genentech, Inc., GlaxoSmithKline, Health Point, Idera, Innocutis, Innovail, Intendis, Isdin, Johnson & Johnson, Laboratory Skin Care Inc., LEO Pharma, L'Oreal, 3M, Maruho, Medical International Technologies, Medicis Pharmaceutical Corp., Merck, Merz, Nano Bio, Novartis AG, Noven Pharmaceuticals, Nucryst Pharmaceuticals Corp., Obagi, Onset, OrthoNeutrogena, PediaPharma, Pfizer, Promius, QLT, Inc., PharmaDerm, PuraCap, Quatrix, Quinnova, Serono (Merck Serono International SA), SkinMedica, Inc., Stiefel Laboratories, Inc., Sun Pharma, Taro, TolerRx, Triax, UCB, Valeant Pharmaceuticals Intl, Warner-Chilcott, XenoPort, and ZAGE. **Z.D. Draelos** received grant support from Arcutis Biotherapeutics, Inc. for the conduct of this study. **S.E. Kempers** is an investigator for Arcutis, and serves as a consultant for Foamix and Kinex. **M. Zirwas** is an investigator, consultant, and/or speaker for AbbVie, Arcutis, Asana, Avillion, Dermavant, DS Biopharma, Edessa Biotech, Fit Bit, Foamix, Galderma, Genentech/Novartis, Incyte, Janssen, L'Oreal, LEO Pharma, Lilly, Menlo, Ortho Dermatologics, Pfizer, Regeneron/Sanofi, and UCB, and is part owner of AsepticMD. **M.-L. Trotman** is a consultant for Arcutis. **K. Smith, D.W. Osborne, L. Navale, C. Merritt, D.R. Berk, and H. Welgus** are employees of Arcutis Biotherapeutics, Inc. **D.W. Osborne and H. Welgus** have a patent application relevant to this work.

This work was supported by Arcutis Biotherapeutics, Inc.

Treatment of Chronic Plaque Psoriasis

Background

- High-potency corticosteroids and vitamin D derivatives are the main treatments, but have long-term tolerability issues
 - Sensitive areas such as the face and intertriginous areas require additional considerations
- PDE-4 activity is elevated in psoriatic skin relative to healthy skin, and inhibition of PDE-4 results in downregulation of inflammatory cytokines TNF α , IFN γ , IL-17, and IL-23^{1,2}
- Roflumilast cream (ARQ-151) is a potent, selective PDE-4 inhibitor under clinical investigation
 - Demonstrates ~25- to >300-fold higher potency than currently available PDE-4 inhibitors³

PDE-4: phosphodiesterase-4.

¹Pincelli C, et al. *J Drugs Dermatol* 2018;17:835-840. ²Li H, et al. *Front Pharmacol* 2018;9:1048. ³Dong C, et al. *J Pharmacol Exp Ther* 2016;358:413-422.


Safety and Efficacy of Topical Once-Daily Roflumilast Cream in Subjects With Chronic Plaque Psoriasis: Objective and Methods

Objective

- Evaluate the safety and efficacy of 2 doses of roflumilast cream versus vehicle

Methods

- Parallel-group, randomized, double-blind, vehicle-controlled Phase 2b Study
- Adult subjects (≥ 18 years) with chronic plaque psoriasis (IGA ≥ 2)
- Roflumilast cream 0.3%, 0.15%, or vehicle; QD for 12 weeks
- Primary endpoint: IGA status of 'clear' or 'almost clear' (score 0 or 1) at Week 6
- Secondary endpoint: For subjects with I-IGA ≥ 2 at baseline, I-IGA success at each time point



ClinicalTrials.gov NCT03638258


Baseline Characteristics

	Roflumilast 0.3% (n=109)	Roflumilast 0.15% (n=113)	Vehicle (n=109)
Age, mean years (SD)	51.7 (14.1)	54.4 (14.2)	55.5 (13.5)
Sex, male, n (%)	56 (51.4)	62 (54.9)	67 (61.5)
Race, n (%)			
White	82 (75.2)	95 (84.1)	92 (84.4)
Black	12 (11.0)	10 (8.8)	7 (6.4)
Multiple/other	15 (13.8)	8 (7.1)	10 (9.2)
Psoriasis-affected BSA, mean % (SD)	6.3 (4.0)	6.4 (3.9)	6.4 (3.6)
IGA score, n (%)			
2 (mild)	17 (15.6)	18 (15.9)	11 (10.1)
3 (moderate)	84 (77.1)	83 (73.5)	89 (81.7)
4 (severe)	8 (7.3)	12 (10.6)	9 (8.3)

	Roflumilast 0.3% (n=109)	Roflumilast 0.15% (n=113)	Vehicle (n=109)
PASI, mean score (SD)	7.7 (3.6)	8.0 (3.9)	7.6 (3.1)
WI-NRS, mean score (SD)	6.1 (2.7)	5.6 (3.1)	5.9 (2.9)
WI-NRS score ≥ 6 , n (%)	71 (65.1)	62 (54.9)	64 (58.7)
PSD, mean total score (SD)	68.9 (41.2)	69.6 (46.2)	75.1 (42.6)
Intertriginous Area			
I-IGA score ≥ 2 , n (%)			
2 (mild)	6 (37.5)	12 (66.7)	7 (41.2)
3 (moderate)	8 (50.0)	3 (16.7)	8 (47.1)
4 (severe)	1 (6.3)	1 (5.6)	2 (11.8)

Data are presented for intent-to-treat population. BSA: body surface area; IGA: Investigator Global Assessment; I-IGA: Intertriginous Investigator Global Assessment; PASI: Psoriasis Area and Severity Index; PSD: Psoriasis Symptom Diary; SD: standard deviation; WI-NRS: Worst Itch Numeric Rating Scale.


Primary Endpoint of IGA 'Clear' or 'Almost Clear' at Week 6 Was Met for Both Roflumilast Cream Doses



Data are presented for intent-to-treat population. CI: confidence interval; IGA: Investigator Global Assessment.


Roflumilast Cream Improved Severity of Chronic Plaque Psoriasis as Measured by IGA Success

Subjects Achieving IGA of 'Clear' or 'Almost Clear' Plus 2-Grade Improvement From Baseline



Data are presented for intent-to-treat population. CI: confidence interval; IGA: Investigator Global Assessment.

Roflumilast Cream Led to Early Improvement in Chronic Plaque Psoriasis Area and Severity Index



* $P < 0.001$ vs vehicle

Data are presented for intent-to-treat population. CI: confidence interval; LS: least squares; PASI: Psoriasis Area and Severity Index.

Roflumilast Cream Improved Severity of Plaque Psoriasis

Roflumilast 0.3%

Roflumilast 0.15%

Vehicle

Baseline

IGA = 3



IGA = 3




IGA = 3



Week 8

IGA = 1



IGA = 1




IGA = 3



IGA: Investigator Global Assessment.

Most Subjects With Intertriginous Plaques Treated With Roflumilast Cream Achieved I-IGA Success by Week 6 With Continued Improvement Through Week 12


Subjects With Intertriginous Plaques Achieving I-IGA of 'Clear' or 'Almost Clear' Plus 2-Grade Improvement From Baseline



Data are presented for intent-to-treat population. CI: confidence interval; I-IGA: Intertriginous Investigator Global Assessment.

Roflumilast Cream Rapidly Improved Subject-Reported Itch


Subjects Achieving 4-Point Reduction in WI-NRS*



*Subjects with score of ≥ 6 on WI-NRS at baseline.

Data are presented for intent-to-treat population. CI: confidence interval; WI-NRS: Worst Itch Numeric Rating Scale.

Roflumilast Cream Rapidly Reduced Patient-Reported Burden of Disease



Data are presented for intent-to-treat population. CI: confidence interval; PSD: Psoriasis Symptom Diary.

Most TEAEs Were Mild or Moderate

- Treatment-related AEs were uncommon and were similar across groups
- More subjects discontinued the study due to an AE in the vehicle group vs the roflumilast groups
- Rates of application site pain were low and similar to vehicle
- 97% of AEs were rated mild or moderate
- Rates of gastrointestinal and psychiatric AEs were low and similar between groups
- Proportion of subjects with weight loss was comparable across groups and comparable to weight gain

TEAE, n (%)	Roflumilast 0.3% (n=109)	Roflumilast 0.15% (n=110)	Vehicle (n=107)
Subjects with any TEAE	42 (38.5)	30 (27.3)	32 (29.9)
Subjects with any treatment-related TEAE	7 (6.4)	3 (2.7)	7 (6.5)
Subjects with any SAE^a	1 (0.9)	1 (0.9)	2 (1.9)
Subjects who discontinued study due to AE^b	1 (0.9)	0	2 (1.9)
Most common TEAE (>2%)			
Upper respiratory tract infection (including viral)	9 (8.2)	8 (7.3)	4 (3.7)
Nasopharyngitis	4 (3.7)	3 (2.7)	4 (3.7)
Application site pain	2 (1.8)	1 (0.9)	3 (2.8)
Sinusitis	3 (2.8)	0	0
Urinary tract infection	0	3 (2.7)	1 (0.9)

^aRoflumilast 0.3%: worsening of chest pain in a subject with history of myocardial infarction; roflumilast 0.15%: melanoma (not in treatment area); vehicle group: acute infarction of left basal ganglia, spontaneous miscarriage. ^bRoflumilast 0.3%: onset of worsening psoriasis; vehicle: mood swings, contact dermatitis.

Conclusions

- PDE-4 inhibition is a validated mechanism of action for oral psoriasis therapy, but new for topical psoriasis treatment
- Roflumilast once-daily cream demonstrated significant improvements in psoriasis signs and symptoms
 - Statistically significant increase in IGA ‘clear’/‘almost clear’ and IGA success
 - Improvement in secondary endpoints of subject-reported itch, burden of disease
 - Significant clinical improvement in intertriginous areas: most patients ‘clear’ at Week 8
- Improvements in efficacy were observed as early as Week 2
- Roflumilast cream was well-tolerated; application site pain similar to vehicle

Roflumilast cream, an investigational once-daily topical PDE-4 inhibitor, achieved improvements in psoriasis signs and symptoms compared with vehicle cream, including in intertriginous areas

Acknowledgements

- **Thank you to the investigators and their staff for their participation in this trial**
 - David Adam, Ajax, Ontario, Canada; Lorne Albrecht, Surrey, British Columbia, Canada; Javier Alonso-Llamazares, Miami, Florida; Suzanne Bruce, Houston, Texas; Laura Ferris, Pittsburgh, Pennsylvania; Rion Forconi, Sanford, Florida; Chih-ho Hong, Surrey, British Columbia, Canada; Terry Jones, College Station, Texas; Mark Lee, San Antonio, Texas; Patricia Lee, Webster, Texas; Wei Jing Loo, London, Ontario, Canada; Charles Lynde, Markham, Ontario, Canada; Vandana Madkan, Santa Monica, California; Walter Nahm, San Diego, California; Lawrence Osman, Northridge, California; David Pariser, Norfolk, Virginia; Sheetal Sapra, Oakville, Ontario, Canada; Daniel Stewart, Clinton Township, Michigan; David Stoll, Beverly Hills, California; Darryl Paul Toth, Windsor, Ontario, Canada; Stephen Tying, Houston, Texas; Marni Charlotte Wiseman, Winnipeg, Manitoba, Canada
- **We are grateful to the study participants and their families for their time and commitment**